



hyperprolinemia

Hyperprolinemia is an excess of a particular protein building block (amino acid), called proline, in the blood. This condition generally occurs when proline is not broken down properly by the body. There are two inherited forms of hyperprolinemia, called type I and type II.

People with hyperprolinemia type I often do not show any symptoms, although they have proline levels in their blood between 3 and 10 times the normal level. Some individuals with hyperprolinemia type I exhibit seizures, intellectual disability, or other neurological or psychiatric problems.

Hyperprolinemia type II results in proline levels in the blood between 10 and 15 times higher than normal, and high levels of a related compound called pyrroline-5-carboxylate. This form of the disorder has signs and symptoms that vary in severity, and is more likely than type I to involve seizures or intellectual disability.

Hyperprolinemia can also occur with other conditions, such as malnutrition or liver disease. In particular, individuals with conditions that cause elevated levels of lactic acid in the blood (lactic acidemia) may have hyperprolinemia as well, because lactic acid inhibits the breakdown of proline.

Frequency

It is difficult to determine the prevalence of hyperprolinemia type I because most people with the condition do not have any symptoms. Hyperprolinemia type II is a rare condition; its prevalence is also unknown.

Genetic Changes

Mutations in the *ALDH4A1* and *PRODH* genes cause hyperprolinemia.

Inherited hyperprolinemia is caused by deficiencies in the enzymes that break down (degrade) proline. Hyperprolinemia type I is caused by a mutation in the *PRODH* gene, which provides instructions for producing the enzyme proline oxidase. This enzyme begins the process of degrading proline by starting the reaction that converts it to pyrroline-5-carboxylate.

Hyperprolinemia type II is caused by a mutation in the *ALDH4A1* gene, which provides instructions for producing the enzyme pyrroline-5-carboxylate dehydrogenase. This enzyme helps to break down the pyrroline-5-carboxylate produced in the previous reaction, converting it to the amino acid glutamate. The conversion between proline and glutamate, and the reverse reaction controlled by different enzymes, are important in

maintaining a supply of the amino acids needed for protein production, and for energy transfer within the cell.

A deficiency of either proline oxidase or pyrroline-5-carboxylate dehydrogenase results in a buildup of proline in the body. A deficiency of the latter enzyme leads to higher levels of proline and a buildup of the intermediate breakdown product pyrroline-5-carboxylate, causing the signs and symptoms of hyperprolinemia type II.

Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. Most often, the parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but do not show signs and symptoms of the condition. In about one-third of cases, individuals carrying one copy of an altered *PRODH* gene have moderately elevated levels of proline in their blood, but these levels do not cause any health problems. Individuals with one altered *ALDH4A1* gene have normal levels of proline in their blood.

Other Names for This Condition

- proline oxidase deficiency
- prolinemia
- pyrroline-5-carboxylate dehydrogenase deficiency
- pyrroline carboxylate dehydrogenase deficiency

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Deficiency of pyrroline-5-carboxylate reductase
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2931835/>
- Genetic Testing Registry: Proline dehydrogenase deficiency
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0268529/>

Other Diagnosis and Management Resources

- Baby's First Test
<http://www.babysfirsttest.org/newborn-screening/conditions/prolinemia>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>

- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Health Topic: Amino Acid Metabolism Disorders
<https://medlineplus.gov/aminoacidmetabolismdisorders.html>
- Health Topic: Genetic Brain Disorders
<https://medlineplus.gov/geneticbraindisorders.html>
- Health Topic: Newborn Screening
<https://medlineplus.gov/newbornscreening.html>

Genetic and Rare Diseases Information Center

- Hyperprolinemia type 2
<https://rarediseases.info.nih.gov/diseases/6710/hyperprolinemia-type-2>

Educational Resources

- Disease InfoSearch: Hyperprolinemia
<http://www.diseaseinfosearch.org/Hyperprolinemia/3600>
- MalaCards: hyperprolinemia
<http://www.malacards.org/card/hyperprolinemia>
- Orphanet: Hyperprolinemia type 1
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=419
- Orphanet: Hyperprolinemia type 2
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=79101
- Screening, Technology and Research in Genetics
<http://www.newbornscreening.info/>

Patient Support and Advocacy Resources

- Children Living with Inherited Metabolic Diseases
<http://www.climb.org.uk/>
- National Organization for Rare Disorders (NORD)
<https://rarediseases.org/rare-diseases/hyperprolinemia-type-i/>
- National Organization for Rare Disorders: Hyperprolinemia Type II
<https://rarediseases.org/rare-diseases/hyperprolinemia-type-ii/>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22hyperprolinemia%22>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28hyperprolinemia%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- HYPERPROLINEMIA, TYPE I
<http://omim.org/entry/239500>
- HYPERPROLINEMIA, TYPE II
<http://omim.org/entry/239510>

Sources for This Summary

- Campbell HD, Webb GC, Young IG. A human homologue of the *Drosophila melanogaster* sluggish-A (proline oxidase) gene maps to 22q11.2, and is a candidate gene for type-I hyperprolinaemia. *Hum Genet.* 1997 Nov;101(1):69-74.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9385373>
- Geraghty MT, Vaughn D, Nicholson AJ, Lin WW, Jimenez-Sanchez G, Obie C, Flynn MP, Valle D, Hu CA. Mutations in the Delta1-pyrroline 5-carboxylate dehydrogenase gene cause type II hyperprolinemia. *Hum Mol Genet.* 1998 Sep;7(9):1411-5.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9700195>
- OMIM: HYPERPROLINEMIA, TYPE I
<http://omim.org/entry/239500>
- OMIM: HYPERPROLINEMIA, TYPE II
<http://omim.org/entry/239510>
- Humbertclaude V, Rivier F, Roubertie A, Echenne B, Bellet H, Vallat C, Morin D. Is hyperprolinemia type I actually a benign trait? Report of a case with severe neurologic involvement and vigabatrin intolerance. *J Child Neurol.* 2001 Aug;16(8):622-3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11510941>

- Jacquet H, Berthelot J, Bonnemains C, Simard G, Saugier-Weber P, Raux G, Campion D, Bonneau D, Frebourg T. The severe form of type I hyperprolinaemia results from homozygous inactivation of the PRODH gene. *J Med Genet*. 2003 Jan;40(1):e7.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12525555>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1735267/>
 - Shivananda, Christopher R, Kumar P. Type I hyperprolinemia. *Indian J Pediatr*. 2000 Jul;67(7):541-3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10957843>
-

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/condition/hyperprolinemia>

Reviewed: June 2007

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services